

More on Lagrangian relaxation

(Wolsey : Chapter 10)

Consider an integer optimization problem of form :

$$(P) \quad z = \max c x$$

$$D x \leq d$$

$$x \in X$$

"complicating constraints"

↪ include the integrality constraints

Let D $m \times n$ matrix

$$d \in \mathbb{R}^m$$

$$x \in \mathbb{R}^n$$

For any $u = (u_1, \dots, u_m) \geq 0$, consider the Lagrangian relaxation :

$$(P_u) \quad z(u) = \max c x + u (d - D x)$$

$$x \in X$$

u : price or dual variables or
Lagrangian multipliers associated
 with $Dx \leq d$

Proposition : (P_u) is a relaxation of (P)
 for all $u \geq 0$.

Proof :

(i) $\{x : Dx \leq d, x \in X\} \subseteq X$

the feasible region of (P_u) is at
 least as large

(ii) $cx + \underbrace{u(d - Dx)}_{\geq 0} \geq cx \quad \forall x \in X$

$\forall x \in X$ the objective value $z_i(P_u)$ is
 at least as great as $z_i(P)$

□

Therefore : $z(u) \geq z \quad \forall u \geq 0$

(maximization problem)

To find the best (\equiv smallest) upper bound $z(u)$ for $u \geq 0$, we solve the lagrangian Dual Problem:

$$(LD) \quad w_{LD} = \min \{ z(u) : u \geq 0 \}$$

Solving a lagrangian relaxation may sometimes lead to solve (P):

let $u \geq 0$:

Proposition: If:

(i) $x(u)$ is an optimal solution to (P_u)

(ii) $Dx(u) \leq d$

(iii) if $u_i > 0$ then $(Dx(u))_i = d_i$
(complementarity)

then $x(u)$ is optimal for (P).

Proof:

From (i):

$$w_{LD} \leq z(u) = c x(u) + u (d - Dx(u))$$